


MythBuster: Eating Disorders

MYTH: "Eating disorders aren't serious – they're just diets gone wrong..."
and other unhelpful myths



Eating Disorders

Eating disorders are among the most serious and misunderstood of all mental disorders. A number of myths and stereotypes exist about eating disorders that can be potentially damaging to young people affected by them and to their families. This MythBuster aims to dispel these myths and present an evidence-based understanding of how eating disorders can affect young people and how their needs can be met.

What are eating disorders?

A person has an eating disorder when their attitudes to food, weight, body size or shape lead to marked changes in their eating or exercise behaviours which interfere with their life and relationships (1). There are three main types of eating disorders: Anorexia Nervosa, Bulimia Nervosa and Eating Disorders Not Otherwise Specified (EDNOS) (2).

The most common type of eating disorder is EDNOS, affecting half of all adolescents and adults diagnosed with an eating disorder (3). A diagnosis of EDNOS is made when a person has symptoms that do not quite fit the criteria for either Anorexia Nervosa or Bulimia Nervosa, but that are seriously interfering with their day-to-day life – for example, affecting their physical health, work or studies or causing problems in their relationships. The most common form of EDNOS is **Binge Eating Disorder**.

Bulimia Nervosa is the next most common eating disorder. It involves an often-secretive cycle of ‘binge’ eating (i.e. eating large amounts of food in a short space of time, accompanied by feeling a loss of control) followed by ‘compensatory’ behaviours (i.e. ‘compensating’ or trying to get rid of the food by vomiting, abusing laxatives, over-exercising or fasting). **These behaviours are usually very distressing to the person as they are experienced as being out of their control and often bring on intense feelings of guilt and shame** (4-5). Bulimia is no less serious when a person uses over-exercising or fasting to try to lose weight, rather than vomiting or abusing laxatives (6-8).

Anorexia Nervosa is the least common of all eating disorders. It involves (a) persistent efforts to lose or maintain a low body weight, (b) a distorted body image (i.e. the way a person sees their body is distorted, making them think they are bigger than they actually are) and (c) an intense fear of gaining weight (2).

The most common myths about eating disorders include:

1. Myths about the seriousness of eating disorders

Eating disorders are often described as being a ‘trend’, ‘phase’, ‘attention-seeking’ or simply ‘a diet gone wrong’. These myths are damaging as they can fuel the belief that people with eating disorders are responsible for their symptoms and could simply ‘stop’ their behaviour if they really wanted to, or tried hard enough (9).

2. Myths about who can be affected by eating disorders

Stereotypes about the ‘types of people’ who develop eating disorders are common (e.g. only females are affected; 9-13). These stereotypes are harmful as they can increase the likelihood that eating disorders will go unnoticed in young people who don’t ‘fit the mould’ (14). They can also make it less likely that these young people will seek help.

3. Myths about treatments for, and recovery from, eating disorders

Beliefs that eating disorders are ‘untreatable’ are still widespread. This adds to a sense of hopelessness that is sometimes associated with eating disorders – not just in the community, but also among health professionals. However, evidence shows that with support and appropriate treatment, most young people recover from their symptoms (3,15-16).

Early treatment is associated with better outcomes (17-19).

Busting the myths...

MYTH: “Eating disorders are a fashion, trend, phase or attention-seeking”

Although some young people might try dieting or other weight-control behaviours because of the influence of their friends, eating disorders don’t develop because a person wants to be fashionable. They are serious illnesses that can severely disrupt every aspect of a person’s life, including their physical health (20-21). **In adolescence, the health risks can be even more serious as an eating disorder can impact on physical growth and development** (22-23).

Similarly, eating disorders are not about ‘attention-seeking’. In fact, people often go to great lengths to hide their behaviours from those around them (11). If someone does notice and approaches them about their weight or behaviour, a person with an eating disorder will often deny that they have a problem (11,22). This may be even more common among adolescents than adults, as adolescents tend to experience greater denial of symptoms and less desire for help (24).

MYTH: “An eating disorder is just a diet gone wrong”

Eating disorders often begin with dieting (9,22-23,25-26), but they are more than just ‘a diet gone wrong’. When a person has an eating disorder they can experience a range of distressing and disabling feelings and behaviours, not just restrictive eating, but bingeing, purging (i.e. vomiting or abusing laxatives) and/or excessive exercising. These behaviours are often experienced as being out of the person’s control. Other mental health difficulties, such as depression and substance use, are also common (see 19,27).

MYTH: “Eating disorders don’t affect males”

Eating disorders are more common among young women than men. However, they also affect males. In adolescence and early adulthood, it’s estimated that 10% of cases of Anorexia Nervosa (28), 10-15% of cases of Bulimia Nervosa (29) and up to 40% of cases of Binge Eating Disorder occur in males (30). Some males with eating disorders may be driven by the desire to lose weight while others may be driven by the desire to gain weight in order to achieve an ideal muscular, trim physique (30).

There is evidence that it takes longer for males to receive professional help and treatment than females (29), which may be explained in part by this myth. Typically it is only when symptoms become severe that a diagnosis of an eating disorder is considered in boys and young men (30). Males may also be even more motivated to hide their symptoms due to the stigma associated with having what they or others perceive to be ‘a female disorder’ (30).

MYTH: “Eating disorders only affect the wealthy and Westerners”

Stereotypes still exist that eating disorders predominantly affect girls from affluent or privileged families (12-13,20,31) and those from Western cultures (e.g. Australia, USA, Europe) where the ‘ideal beauty’ involves being thin and toned (9,11-12,19). However evidence shows that **anyone can develop an eating disorder, regardless of ethnicity or social class** (9,12-13,20,22,31).

MYTH: “It’s easy to tell if someone has an eating disorder because they will be very underweight”

You can’t tell just by looking at someone whether or not they have an eating disorder. A person with an eating disorder may be underweight, within a normal weight range, or overweight (2).

MYTH: “Families are to blame for eating disorders”

A particularly harmful myth is that eating disorders only occur in certain types of families and are ‘caused’ by certain parenting styles (18). In the past, parents were often prevented from having any involvement in their child’s treatment as they were seen as ‘part of the problem’ (18). Unfortunately, many parents still report feeling ‘blamed’ for their child’s eating disorder (e.g. 32). There is no evidence to suggest that eating disorders are caused by particular parenting styles (11,18,33). **Parents and families should be seen as part of the solution to overcoming an eating disorder, not the problem.** Clinical best-practice guidelines state that family members should normally be included in the treatment of adolescents with eating disorders (34).

MYTH: “The solution to eating disorders is simple – just stop”

Supporting someone with an eating disorder can sometimes be challenging. It can be difficult to understand why the person can’t change their thoughts about themselves or stop their behaviour and return to ‘normal’. Unfortunately, recovery is not as simple as ‘just stopping’. People with eating disorders need to be supported to learn how to think more realistically about their body and more positively about themselves, and to learn techniques for managing difficult emotions without turning to food or weight-control behaviours.

Busting the myths on weight loss...

As well as busting the myths surrounding eating disorders in general, it is important to tackle some unhelpful myths about ‘what works’ for weight loss and what effect different weight loss strategies can have on your health. People may believe that unhealthy weight loss strategies work and/or that they’re not really a serious problem (35). This is a myth, as evidence shows that not only are many of these techniques ineffective ways to lose weight, but they can be very damaging to the person’s physical and psychological health (4-8,36).

MYTH: “There is no such thing as too much exercise”

Exercising excessively is not good for you physically or emotionally. Over-exercising can cause serious health problems, including osteoporosis (brittle bones), lowered hormones, heart problems and permanent damage to joints and tendons (37). It can also lead to strong feelings of guilt and depressed mood when exercising goals are not met, or when a person is unable to exercise (38).

MYTH: “Vomiting or using laxatives gets rid of the calories consumed by eating”

Even when done immediately after eating, vomiting does not get rid of all of the calories a person has consumed. By the time food reaches the stomach, most of the calories have already been absorbed by the body. Similarly, laxatives don’t prevent calories being absorbed, so they have almost no impact on weight loss. Both chronic vomiting and laxative abuse can cause serious health problems that can be life-threatening (39-40). Chronic vomiting can cause tearing or bleeding in the oesophagus (the ‘food pipe’), digestive problems and dental damage. Abusing laxatives can cause serious problems with bowel functioning, such as bloating, gas, pain and loss of control over bowel movements (41). Both of these behaviours can cause electrolyte imbalances that can be life-threatening (41-42).

MYTH: “Fad dieting is a good way to lose weight”

Fad diets (or any short-term weight loss strategy) might work in the short-term, but in the long-run, they usually lead to weight gain, not loss (36). This is because dieting slows down your body’s metabolism making it harder to burn-off calories consumed. Dieting also increases a person’s pre-occupation with food and craving, **such that they may actually end up eating more than they would if they were not dieting.**

So what does this all mean?

- Eating disorders are serious, damaging and potentially life-threatening illnesses – they need to be taken seriously, not trivialised.
- There are many unhelpful stereotypes about people with eating disorders. We need to be aware of these stereotypes and challenge them wherever possible.
- Eating disorders are complex conditions that are caused by a combination of factors. It is misguided to blame someone for an eating disorder - either the individual affected or their family. Eating disorders are not a lifestyle choice.
- **Recognizing the symptoms or warning signs of eating disorders is critical, as the earlier a problem is detected and treated, the better the outcome for the young person** (see the **headspace** factsheet on Eating Disorders for more information on early warning signs – www.headspace.org.au). Early treatment increases the chances of fully recovering, and recovering faster.

Getting help

Worried that you may be at-risk of developing, or have, an eating disorder?

It is strongly recommended that you consult a health professional if you feel you may have symptoms of an eating disorder. In addition to professional help, there is a useful online program called ‘*Overcoming Disordered Eating*’ (download at <http://www.cci.health.wa.gov.au/resources/consumers.cfm>). There are other self-help resources available (e.g. ‘Getting Better Bit(e) by Bit(e): A survival kit for sufferers of Bulimia Nervosa and Binge Eating Disorder’; 43), however it’s not recommended to try to treat an eating disorder using self-help alone – getting professional help as early as possible is important. The best place to start is to contact your GP or your nearest headspace centre (www.headspace.org.au).

Worried about someone who may have/be developing an eating disorder?

It’s important that you approach the person about your concerns and seek advice and support from a health professional. *The Mental Health First Aid Guidelines on Eating Disorders* provide helpful evidence-based advice on the best steps to follow in approaching the person about your concerns, encouraging them to get professional help and looking after yourself (www.mhfa.com.au). You can also call the Butterfly Foundation’s helpline 1800 334 673 to speak to somebody about your concerns or email: support@thebutterflyfoundation.org.au.

Even if the young person denies that they have a problem, it is very important to seek help from a health professional about your concerns. The best place to start is to contact your GP or your nearest headspace centre (www.headspace.org.au). Sometimes, the process of looking for help can be frustrating, it’s important to be persistent in your efforts. There is help out there.

Supporting somebody who has an eating disorder?

Resources that may be helpful are available on The Butterfly Foundation’s website (www.thebutterflyfoundation.org.au) and the Victorian Centre of Excellence in Eating Disorders’ website (www.ceed.org.au).

- Mental Health First Aid Training and Research Program (2008) *Eating disorders: first aid guidelines for assisting adults*. Melbourne: Orygen Youth Health Research Centre, University of Melbourne.
- American Psychiatric Association (2000) *Diagnostic and statistical manual of mental disorders, fourth edition, text revision (DSM-IV-TR ©)*
- Fairburn CG, & Cooper Z (2007) Thinking afresh about the classification of eating disorders. *Int J Eat Disord*, 40: p.107-10.
- Patton GC, Carlin JB, Shao Q, Hibbert ME, Rosier M, Selzer R et al. (1997) Adolescent dieting: healthy weight control or borderline eating disorder? *J Child Psychol & Psychiatry*, 38: p.299-306
- Shisslak CM, Crago M & Estes LS (1995) The spectrum of eating disturbances. *Int J Eat Disord*, 25: p.209-19
- Mond JM, Hay PJ, Rodgers B, Owen C & Mitchell JE (2006) Correlates of the use of purging and non-purging methods of weight control in a community sample of women. *Aust NZ J Psychiatry*, 40: p.136-42
- Mond JM, Hay PJ, Rodgers B, Owen C, Crosby R & Mitchell JE (2006) Use of extreme weight control behaviors with and without binge eating in a community sample of women: implications for the classification of bulimic eating disorders. *Int J Eat Disord*, 39: p.294-302
- Mond JM, Hay PJ, Rodgers B, Owen C & Beumont PJV (2004) Beliefs of women concerning the severity and prevalence of bulimia nervosa. *Int J Eat Disord*, 39: p.299-304
- Striegel-Moore RH & Bulik CM (2007) Risk factors for eating disorders. *Am Psychol*, 62(3): p.181-98
- Bruch H (1978) *The golden cage*. Cambridge, MA: Harvard University Press, 1978
- Polivy J & Herman C (2002) Causes of eating disorders. *Annu Rev Psychol*, 53: p.187-213
- Soh NL, Touyz W & Surgenor LJ (2006) Eating and body image disturbances across cultures: a review. *Eur Eat Disorders Review*, 14: p.54-65
- Gard MCE & Freeman CP (1996) The dismantling of a myth: a review of eating disorders and socioeconomic status. *Int J Eat Disord*, 20(1):p.1-12
- Dolan BM, Lacey JH & Evans C (1990) Eating behaviour and attitudes to weight and shape in British women from three ethnic groups. *Brit J Psychiatry*, 157: p.523-28
- Strober M, Freeman R & Morrell (1997) The long-term course of severe anorexia nervosa in adolescents: survival analysis of recovery, relapse and outcome predictors over 10-15 years in a prospective study. *Int J Eat Disord*, 22(4): p.339-60.
- Steinhausen HC, Botadjeva S, Grigoriou-Serbanescu M, Seidel R & Winkler Metzke C (2000) A transcultural outcome study of adolescent eating disorders. *Acta Psychiatr Scand*, 101: p.60-66
- Bryant-Waugh, R (1993). *Prognosis and outcome*. In B. Lask & R. Bryant-Waugh (Eds.), *Childhood onset anorexia nervosa and related eating disorders* (pp.91-108). Hove: Lawrence Erlbaum.
- Powers PS & Santan CA (2002) Childhood and adolescent anorexia nervosa. *Child Adolesc Psychiatr Clin N Am*, 11: p.219-35
- Reijonen JH, Pratt HD, Patel DR & Greydanus DE (2003) Eating disorders in the adolescent population: an overview. *J Adol Res*, 18(3): p.209-222
- Bulik CM (2001) Eating disorders in adolescents and young adults. *Child Adolesc Psychiatr Clin N Am*, 11: p.201-18
- Katzman DK (2005) Medical complications in adolescents with anorexia nervosa: a review of the literature. *Int J Eat Disord*, 37(S1): p.S52-59
- Herpetz-Dahlman B (2008) Adolescent eating disorders: definitions, symptomatology and comorbidity. *Child Adolesc Psychiatr Clin N Am*, 18: p.31-47
- Steiner H & Lock J (1998) Anorexia nervosa and bulimia nervosa in children and adolescents: a review of the past 10 years. *J Am Acad Child Adolesc Psychiatry*, 37: p.353-59
- Fisher M, Schneider M, Burns J, Symons H & Mandel FS (2001) Differences between adolescents and young adults at presentation to an eating disorders program. *J Adol Health*, 28(3): p.222-27
- French SA & Jeffrey RW (1994) Consequences of dieting to lose weight: Effects on physical and mental health. *Health Psychology*, 13(3): p.195-212
- Brewerton TD, Dansky BS, Kilpatrick DG & O'Neil PM (2000) Which comes first in the pathogenesis of bulimia nervosa, dieting or bingeing? *Int J Eat Disord*, 28: p.259-64
- LeGrange D & Loeb KL (2007) Early identification of eating disorders: prodrome to syndrome. *Early Intervention in Psychiatry*, 1: p.27-39
- Hsu LKK (1994) Epidemiology of the eating disorders. *Psychiatr Clin North Am*, 19: p.681-700
- Carlat DJ, Camargo CA & Herzog DB (1997) Eating disorders in males: a report of 135 patients. *Am J Psychiatry*, 154: p.1127-32
- Muise AM, Stein DG & Gordon A (2003) Eating disorders in adolescent boys: a review of the adolescent and young adult literature. *J Adol Health*, 33: p.427-35
- Wildes JE, Emery RE & Simons AD (2001) The role of ethnicity and culture in the development of eating disturbance and body dissatisfaction: a meta-analytic review. *Clin Psychol Rev*, 21(4): p.521-555
- Beat (2008) *Failing Families*. Norwich: Beat
- The Royal Australian and New Zealand College of Psychiatrists (2009) *Anorexia Nervosa: Australian treatment guide for consumers and carers*. Melbourne: RANZCP
- National Institute of Clinical Excellence (2004) *Clinical Guideline 9: Eating disorders - core interventions in the treatment and management of anorexia nervosa, bulimia nervosa and related eating disorders*. London: NICE
- Mond JM (2006) Self-recognition of disordered eating among women with bulimic-type eating disorders: a community-based study. *Int J Eat Disorders*, 39(8): p.747-53
- Field AE, Austin SB, Taylor CB et al. (2001) Relation between dieting and weight change among pre-adolescents and adolescents. *Pediatrics*, 112(4): p.900-6.
- Adams J & Kirby R (2001) Exercise dependence and overtraining: the physiological and psychological consequences of excessive exercise. *Res Sports Med*, 10(3): p.199-222
- Mond JM & Calegero RM (2009) Excessive exercise in eating disorder patients and in healthy women. *Aust N Z J Psychiatry*, 43: p.227-234
- Tylka TL & Subich LM (2002) Exploring young women's perceptions of the effectiveness and safety of maladaptive weight control techniques. *J Couns Dev*, 80(1): p.101-110
- Keel, P. K. (2007) Purging disorder: Subthreshold variant or full-threshold eating disorder? *Int J Eat Disord*, 40: p.589-94.
- Neims DM, McNeill J, Giles TR, et al. (1995) Incidence of laxative abuse in community and bulimic populations: A descriptive review. *Int J Eat Disord*, 17(3): p.211-28
- Turner J, Battik M, Palmer LJ, Forbes B & McDermott BM (2000) Detection and importance of laxative use in adolescents with anorexia nervosa. *J Amer Acad Child & Adolesc Psychiatry*, 39(3): p.378-38
- Schmidt U & Treasure J (1993) *Getting Better Bit(e) by Bit(e): A survival kit for sufferers of Bulimia Nervosa and Binge Eating Disorder*. East Sussex: Psychology Press

Acknowledgements

headspace MythBusters are prepared by the Centre of Excellence in Youth Mental Health. The series aims to unveil common myths that are contrary to the research evidence about mental health and substance use problems affecting young people. Experts on the topic have reviewed the summary before publication, including young people. The authors would like to thank all of the experts consulted for their input.

MythBuster Writers
Ms Faye Scanlan
Assoc Prof Rosemary Purcell

Clinical Consultants
Prof. Susan Paxton
Ms Laura Hart
Ms Michelle Robertson
Ms Claire Diffey
Ms Carmen Garret

headspace National Youth Mental Health Foundation Ltd is funded by the Australian Government Department of Health and Ageing under the Youth Mental Health Initiative Program.

For more details about **headspace** visit www.headspace.org.au.

Copyright © 2010 Orygen Youth Health Research Centre
This work is copyrighted. Apart from any use permitted under the Copyright Act 1968, no part may be reproduced without prior written permission from Orygen Youth Health Research Centre.
ISBN: 978-0-9807780-2-1 (Online): 978-0-9807780-3-8.